


Appl. No. Unassigned; Docket No. NL02 0571 US  
Amdt. dated 01-DEC-2005  
Preliminary Amendment

Amendments to the Claims

1. (ORIGINAL) A phenylethanediol derivative, characterized in that the phenylethanediol derivative comprises at least one photo-convertible group suitable for adjusting the helical twisting power of the phenylethanediol derivative.
2. (ORIGINAL) The phenylethanediol derivative of claim 1 further having at least one polymerizable group.
3. (CURRENTLY AMENDED) The phenylethanediol derivative of ~~claim 1~~  
~~or 2~~claim 1 wherein the photo-convertible group is a photo-isomerizable group.
4. (ORIGINAL) The phenylethanediol derivative of claim 3 wherein the photo-isomerizable group is an olephinic group.
5. (CURRENTLY AMENDED) The phenylethanediol derivative of ~~any one~~  
~~of claims 1-4~~claim 1 wherein the polymerizable group is a (meth)acrylate group.
6. (CURRENTLY AMENDED) The phenylethanediol derivatives of ~~any one~~  
~~of the preceding claims~~claim 1 wherein the phenylethanediol has the formula

Appl. No. Unassigned; Docket No. NL02 0571 US  
 Amdt. dated 01-DEC-2005  
 Preliminary Amendment



wherein

A stands for a bond or a p-phenylene group;

B and B' are independently  $(O)_p-C_oH_{2o}-O-CO-CR'=CH_2$ , o being 2-12, p being 0 or 1, and R' being H or  $CH_3$ ;

P stands for a  $CH_2$  or a  $C=O$  group;

Q and Q' are independently selected from H, C1-C3 alkyl, C1-C3 alkoxy, halogen, and CN;


n is an integer from 1 to 3; and

m is an integer from 0 to 2.

7. (CURRENTLY AMENDED)

The phenylethanediol derivative of any one of the preceding claims claim 1 wherein the phenylethanediol has the formula

Appl. No. Unassigned; Docket No. NL02 0571 US  
 Amdt. dated 01-DEC-2005  
 Preliminary Amendment



wherein

A stands for a bond or a p-phenylene group;

B is  $(O)_p-C_6H_{2p-1}-O-CO-CR'=CH_2$ ,  $\circ$  being 2-12, p is 1, and R' being H or  $CH_3$ ;

P stands for a  $CH_2$  or a  $C=O$  group;

Q is selected from H, C1-C3 alkyl, C1-C3 alkoxy, halogen, and CN; and

m is an integer from 0 to 2.

**8. (ORIGINAL)** A method for the preparation of the phenylethanediol derivative of claim 1 by the steps of a) synthesizing a 2-hydroxy ether-protected phenylethanediol, b) followed by etherification or esterification of the 1-hydroxy group of the 2-hydroxy ether-protected phenylethanediol with an alcohol (or derivative thereof) or acid, respectively, optionally comprising polymerizable and/or photo-convertible groups, c) then cleaving the ether-protective group to obtain a phenylethanediol derivative with a free 2-hydroxy group, and optionally d) esterification of the free 2-hydroxy group with an acid which optionally comprises one or more polymerizable and/or photo-convertible groups.

**9. (CURRENTLY AMENDED)** A cholesteric composition comprising the phenylethanediol derivative of any one of claims 1-7claim 1.

**10. (CURRENTLY AMENDED)** An optical element, preferably an optical color filter, comprising the phenylethanediol derivative of any one of claims 1-7.

**11. (CURRENTLY AMENDED)** Use of the phenylethanediol derivative of any one of claims 1-7claim 1 in optical elements.